

A **sprain** is an injury that causes a stretch or tear of one or more ligaments in a joint. Ligaments are strong bands of tissue that connect bones at the joint. Sprains may be classified as mild, moderate, or severe. The most common type of sprain involves the ligaments on the outside part of the ankle (lateral ankle sprain). Ligaments on the inside of the ankle may also be injured (medial ankle sprain) as well as ligaments that are high and in the middle of the ankle (high ankle sprains).

An ankle sprain is caused by twisting your ankle. Your foot usually turns in or under but may turn to the outside. Symptoms include mild aching to sudden pain; swelling; discoloration; inability to move the ankle properly; pain in the ankle even when you are not putting any weight on it.

To diagnose a sprained ankle, the healthcare provider will review how the injury occurred and consider your symptoms. He/she will examine your ankle carefully. X-rays may be taken of your ankle.

Treatment:

- Applying ice packs to your ankle for 20 to 30 minutes every 3 to 4 hours for the first 2 to 3 days or until the pain goes away. Thereafter, ice your ankle at least once a day until the other symptoms are gone.
- Elevating your ankle by placing a pillow underneath your foot. Try to keep your ankle above the level of your heart.
- Wrapping an elastic bandage around your ankle to keep the swelling from getting worse.
- Wearing a lace-up brace or ankle stirrup (an Aircast or Gel cast).
- Using crutches until you can walk without pain.
- Taking anti-inflammatory medication, such as ibuprofen, or other pain medication prescribed by your provider. Adults aged 65 years and older should not take non-steroidal anti-inflammatory medicine for more than 7 days without their healthcare provider's approval.
- Doing ankle exercises to improve your ankle strength and range of motion. The exercises will help you return to your normal activity or sports.
- Rarely, severe ankle sprains with complete tearing of the ligaments need surgery. After surgery your ankle will be in a cast for 4 to 8 weeks.

The length of recovery depends on many factors such as your age, health, and if you have had a previous ankle injury. Recovery time also depends on the severity of the sprain. A mild ankle sprain may recover within a few weeks, whereas a severe ankle sprain may take 6 weeks or longer to recover. Recovery also depends on which ligaments were torn. A lateral sprain (outside ligaments) takes less time to recover than a medial sprain (inside ligaments) or a high ankle sprain (high, middle ligaments).

Everyone recovers from an injury at a different rate. Return to your activities will be determined by how soon your ankle recovers, not by how many days or weeks it has been since your injury has occurred. In general, the longer you have symptoms before you start treatment, the longer it

will take to get better. The goal of rehabilitation is to return you to your normal activities as soon as is safely possible. If you return too soon you may worsen your injury.

To help prevent an ankle sprain, wear proper, well-fitting shoes when you exercise. Stretch gently and adequately before and after athletic or recreational activities. Avoid sharp turns and quick changes in direction and movement. Consider taping the ankle or wearing a brace for strenuous sports, especially if you have a previous injury.